arXiv:1405.4796v1 [astro-ph.CO] 19 May 2014

Extremely large peculiar motion of the solar system detected using redshift distribution of distant quasars

Ashok K. Singal*

Astronomy and Astrophysics Division, Physical Research Laboratory,

Navrangpura, Ahmedabad - 380 009, India

(Dated: May 20, 2014)

According to the cosmological principle, an observer fixed in the co-moving co-ordinate system of the expanding Universe should find the Universe to be isotropic, without any preferred directions. To such an observer the redshift distribution due to Hubble expansion of the Universe should also appear similar in all directions. However a peculiar motion of such an observer could introduce a dipole anisotropy in the observed redshift distribution; in reverse an observed dipole anisotropy in the observed redshift distribution could be used to infer the peculiar velocity of the observer with respect to the average Universe. We determine the peculiar velocity of the solar system relative to the frame of distant quasars, by studying the dipole anisotropy, if any, in the redshift distribution of a large sample of quasars distributed across the sky. The magnitude of the peculiar velocity thus determined turns out be extremely large (9750 ± 550 km/s; $\sim 3\%$ the speed of light), and is about an order of magnitude larger than the velocity determined from the dipole anisotropy in the Cosmic Microwave Background Radiation or the value determined earlier relative to the frame of distant radio sources. Even the direction of the motion is in a direction nearly opposite to the earlier determinations. The large differences in the magnitudes of inferred motion as well as their opposite signs are rather disconcerting. A genuine difference between these velocity vectors would imply highly anisotropic Universe, with anisotropy changing with epoch. This would violate the cosmological principle where the isotropy of the Universe is assumed for all epochs, and which is the basis of modern cosmological models.

The peculiar velocity of the solar system relative to the frame of reference provided by the Cosmic Microwave Background Radiation (CMBR) was determined by COBE and WMAP

satellites [1,2] to be 369 km s^{-1} in the direction RA= 168°, Dec= −7°. Though an early work using number counts of radio sources [3] from the NRAO VLA Sky Survey (NVSS) dataset [4] comprising 1.8 million sources, had found the velocity to be consistent (within errors) with that from the CMBR, but a later study [5] using both number counts and the sky brightness arising from discrete radio sources in the same sample found the solar motion to be 1600 ± 400 km/s, about 4 times larger than the CMBR value, though the direction (RA= $157° \pm 9°$, Dec= $-3° \pm 8°$) turned out to be in agreement with the CMBR value. Subsequently two independent groups [6,7], used the same NVSS sample, with one of them additionally using the Westerbork Northern Sky Survey (WENSS) [8], to critically reexamine the two conflicting results and they both found speeds $\sim$ 4 times higher than the CMBR value, corroborating the claims of [5]. This large difference in the CMBR and NVSS speeds is rather puzzling, more so as the direction of the motion turns out to be approximately the same in both cases. Recently there have even been speculations whether the observed excess radio dipole could be an impact of some structure [9] in radio source distribution, though at much larger scale than the local super clusters of optical galaxies. An understanding of the observed difference in speed values is crucial as it relates to the cosmological principle. Therefore it is important if some independent confirmation of the previous results could be made.

In order to resolve this germane issue of two conflicting magnitudes of solar motion inferred from two different methods, we have attempted to determine here in an independent and rather more direct method, our peculiar motion from the anisotropy, if any, in the redshift distribution of a large sample of quasars on the sky. Our sample comprises DR10Q catalogue [10] from the Baryon Oscillation Spectroscopic Survey (BOSS) of the Sloan Digital Sky Survey (SDSS) III. The catalogue contains 166,583 quasars, detected over 6,373 deg^2 covering different regions of the sky, with robust identifications and redshifts measured by a combination of principal component eigenspectra, along with a visual inspection of their spectra. Since the redshift distribution will provide a direct estimate of the peculiar velocity with respect to the reference frame of the quasars, it would also obviate the need for looking for structure formations on large scales that might masquerade as a dipole asymmetry purportedly due to observer's motion, in other indirect methods.

An observer moving with a peculiar velocity $\mathbf{v}$, will find a quasar lying at an angle θ (as

seen in observer's frame) with respect to the direction of motion, to have a redshift,

$$(1+z) = (1+z_{\rm o})/\delta \approx (1+z_{\rm o})(1-(v/c)\cos\theta) \tag{1}$$

where z is the measured redshift of the quasar, $z_{\rm o}$ is the cosmological redshift of the quasar with respect to the comoving rest frame at observer's location, and $\delta = [\gamma(1-(v/c)\cos\theta]^{-1}$ is the Doppler factor due to the observer's motion with $\gamma = 1/\sqrt{(1-(v/c)^2}$ as the Lorentz factor and c as the speed of light in vacuum. Here we shall use the non-relativistic formula for the Doppler factor as both CMBR and NVSS results indicated that $v \ll c$. The observed redshift (Eq. (1)) then has a dipole component $\Delta z = z - z_{\rm o} = -(1+z_{\rm o})(v/c)\cos\theta$, arising from the peculiar motion of the sun with respect to the comoving rest frame. The distribution of the cosmologicl redshift $z_{\rm o}$ should to be statistically independent of the direction (due to the assumed isotropy of the Universe, à la cosmological principle), which implies that if an average of the observed redshift z is taken over all directions in the sky, $\bar{z} = \bar{z}_{\rm o}$, with the dipole component getting cancelled.

Since the sample does not have a uniform sky coverage [10], the direction of the dipole could not be determined as in [5] or in a more formal way using spherical harmonics. However if we already have some idea of the direction of the dipole from other considerations (viz. the dipole direction seen in CMBR or in the NVSS sample), then we can calculate the magnitude of the redshift dipole. For every quasar position in the sky, first we calculate the polar angle θ with respect to the assumed dipole direction. Then a plot of the observed redshifts z against $\cos\theta$ for the sample should yield a linear correlation, $z = a + b\cos\theta$, with $v/c = -b/(1+a)$ giving us the magnitude of the peculiar solar motion.

We have this way determined the magnitude of the dipole in the direction of the NVSS radio dipole (RA= 157°, Dec= −3°) [5] by making a least square fit to the $(\cos\theta, z)$ data on 166583 quasars. It gave an average redshift $\bar{z} = a = 2.209$ and a finite value of the redshift dipole, presumably representing peculiar solar motion, yielding a velocity magnitude of -6620 ± 280 km/s in this direction. In order to ensure that this dipole is not merely due to a skew distribution with a small number of extreme values, we divided the sky into 6 slices of 30° wide bands in polar angle and for each i^{th} band determined the mean redshift $\bar{z}_i$ and its rms due to the spread in redshift within the band. These values are plotted in Fig. 1, and as can be seen from the figure, there is a systematic linear variation in the estimated solar motion component with $\cos\theta$, matching with the straight line fit to the whole data shown

as a dotted line. The computed quantities during each step for various bins are presented in Table 1, which is mostly self-explanatory. Redshift dipole Δz in column (5) is obtained by subtracting the mean value of the redshift over the whole sample, $\bar{z} = \bar{z}_{\rm o} = 2.209$ from the average redshift $\bar{z}_{\rm i}$ in column (4). Column (6) gives the velocity dipole component $(v/c)\,\overline{\cos\theta} = -\Delta z/(1+\bar{z}_{\rm o})$ in units of (10^{-2}) while the calculated peculiar solar velocity component in units of (10^3 km s^{-1}) in the direction of the bin is given in column (7). We get a similar velocity magnitude of -6870 ± 280 km/s in the direction (RA= 168°, Dec= $-7°$) of CMBR dipole.

Intrigued by these unexpectedly large and negative speed values towards the NVSS and CMBR dipole directions, we decided to do a more thorough investigation of the velocity vector by dividing the sky into regions of $10° \times 10°$, covering the whole sky in more than 400 grids, and determined the component of the solar motion for each grid direction following the procedure described above. Thence determined solar motion as a function of the sky position is shown in a 3d plot in Fig. 2. One thing is quite clear that it is not any random fluctuation giving large peculiar motion in some direction. There is a systematic variation in the redshift over the sky which is reflected in the inferred velocity component of the solar motion for various grid points. The plot also shows clearly large negative velocity components near the CMBR or NVSS dipole directions.

To test our procedure as well as to rule out any subtle effects, if any, due to the incomplete sky coverage, we did Monte Carlo trials, for the given quasar positions (thereby keeping the same sky coverage) but with a random redistribution of the redshifts observed in the sample. Five hundred different trials were made, in no case any significant solar motion was found in any of the grid points on the sky. Then an artificial dipole due to a hypothetical solar motion of a different magnitude and direction was superimposed on each of the 500 randomly redistributed redshift data above, and our procedure was repeated to see if we do retrieve the values that we had put in. Not only did we recover that input values, both in direction and magnitude, we also found that the simulated data gave 3d plots very much resembling the one in Fig. 2.

About the grid point that might be close to the apex of the solar motion, one should see a $\cos\theta$ dependence in the inferred velocity component for various other grid points as one moves away from that direction. To determine that, with respect to every grid point in turn on the sky, we computed $\cos\theta$ for all other grid points, and made a least squares fit to a

straight line to the $\cos\theta$ – velocity component data for all these grid points. The best fit with least chi-square value gave us the peculiar solar motion to be 9750 ± 550 km s^{-1} in the direction RA$= 0^\circ \pm 15^\circ$, Dec$= 30^\circ \pm 20^\circ$. Fig. 3 shows the plot for this grid position. Like earlier we also divided the sky into a number of bins for different θ ranges and calculated the average speed for each of these bins. The fit to a straight line is excellent.

It should be noted that the antipole direction (RA$= 180^\circ \pm 15^\circ$, Dec$= -30^\circ \pm 20^\circ$) is approximately within one sigma uncertainty of the CMBR or NVSS dipole directions, hence the solar motion from the quasar redshifts appears to be nearly in a direction opposite to that inferred from CMBR or NVSS. This implies that there is certainly some peculiarity along this direction in sky. However vast differences in the inferred motion (~ 6 times the NVSS value and as much as a factor of ~ 25 or so larger than the CMBR value) cannot be easily explained. A genuine discrepancy in the three dipoles inferred with respect to three different reference frames, would imply a relative motion between the three frames, not in accordance with the present ideas of cosmology.

---

* Electronic address: asingal@prl.res.in

[1] Lineweaver, C. H. et al. *Astrophy. J.* **470,** 38-42 (1996).

[2] Hinshaw, G. et al. *Astrophy. J. Supp. Ser.* **180,** 225 (2009).

[3] Blake, C. & Wall J. *Nature* **416,** 150-152 (2002).

[4] Condon, J. J. et al. *Astr. J.* **115,** 1693-1716 (1998).

[5] Singal, A. K. *Astrophys. J.* **742**, L23-26 (2011).

[6] Rubart, M. & Schwarz, D. J. *Astr. and Astrophys.* **555**, A117 (2013).

[7] Kothari, R., Naskar, A., Tiwari, P., Nadkarni-Ghosh, S. & Jain, P. *preprint* arXiv:1307.1947v2 (2013).

[8] Rengelink, R. B., Tang, Y., de Bruyn, A. G., Miley, G. K., Bremer, M. N., Roettgering, H. J. A. & Bremer, M. A. R. *Astr. and Astrophys. Supp. Ser.* **124**, 259-280 (1997)

[9] Rubart, M., Bacon, D. & Schwarz, D. J. *preprint* arXiv:1402.0376v1 (2014).

[10] Paris, I. et al. *Astr. and Astrophys.* **in press** arXiv:1311.4870v1 (2014).

TABLE I: Velocity components derived from the mean redshifts for different polar angle bins about the direction RA= 157°, Dec= −3°.

Polar-angle range	N i^{th} bin	Mean $\overline{\cos\theta_{\rm i}}$	Bin mean $\bar{z}_{\rm i}$	Redshift dipole $\Delta z_{\rm i} = \bar{z}_{\rm i} - \bar{z}_{\rm o}$	Velocity dipole $(-\Delta z_{\rm i}/1+\bar{z}_{\rm o})(10^{-2})$	Velocity component $v\,\overline{\cos\theta_{\rm i}}(10^3$ km s$^{-1})$
(1)	(2)	(3)	(4)	(5)	(6)	(7)
$0° < \theta_1 \leq 30°$	30233	0.94	2.296	0.087 ± 0.004	-2.71 ± 0.13	-8.1 ± 0.4
$30° < \theta_2 \leq 60°$	54925	0.69	2.246	0.037 ± 0.003	-1.15 ± 0.10	-3.4 ± 0.3
$60° < \theta_3 \leq 90°$	37031	0.28	2.220	0.011 ± 0.004	-0.35 ± 0.13	-1.0 ± 0.4
$90° < \theta_4 \leq 120°$	11072	−0.17	2.229	0.020 ± 0.007	-0.63 ± 0.23	-1.9 ± 0.7
$120° < \theta_5 \leq 150°$	15658	−0.69	2.159	-0.050 ± 0.007	1.56 ± 0.20	4.7 ± 0.6
$150° < \theta_6 \leq 180°$	17664	−0.96	2.142	-0.067 ± 0.006	2.09 ± 0.19	6.3 ± 0.6



FIG. 1: Variation of the inferred velocity component of the solar motion against $\cos\theta$, were θ is the polar angle with respect to the assumed direction of motion (RA= 157°, Dec= −3°) of NVSS dipole. Larger empty circles represent bin-average of the velocity component for each of the 6 different slices of the sky. The dotted line, with small error bars at $\cos\theta = \pm 0.9$, shows the least square fit to straight line for the whole sample of 166583 quasars.



FIG. 2: A 3-d plot of the distribution of the component of the peculiar velocity estimated in the direction of the centre of each $10^\circ \times 10^\circ$ grid, the whole sky covered with > 400 grids.



FIG. 3: Velocity component against $\cos\theta$, where θ is the polar angle with respect to the inferred apex (RA= 0°, Dec= 30°) of the peculiar solar motion. Larger empty circles show average value of velocity component for 10 different slices of the sky, joined by dashed lines. The dotted line represents a least square fit to a straight line to the whole data.